

The quantum black hole in 2+1 dimensions

B. RAM^{a,b}, J. SHIRLEY^b

^aPhysics Department, New Mexico State University
Las Cruces, NM 88003, USA

^bPrabhu-Umrao Institute of Fundamental Research
A2/214 Janak Puri, New Delhi 110 058, India

Abstract

In this paper we investigate the quantum nature of a 2+1 dimensional black hole using the method [arXiv: gr-qc/0504030] which earlier revealed the quantum nature of a black hole in 3+1 dimensions.

The classical analog in 2+1 dimensional Einstein theory with a negative cosmological constant $\Lambda = \frac{-1}{\ell^2}$ of the Schwarzschild line element was first obtained by Bañados, Teitelboim and Zanelli (BTZ) [1]. It is given by [2,3]

$$ds^2 = \left(-8M + \frac{r^2}{\ell^2}\right) dt^2 - \left(\frac{dr^2}{\left(-8M + \frac{r^2}{\ell^2}\right)} + r^2 d\theta^2\right), \quad (1)$$

M being the Arnowitt-Deser-Misner (ADM) mass. Whereas the Schwarzschild line element is asymptotically flat, (1) is not.

The gravitational collapse of a disc of pressureless dust (the 2+1 dimensional analog of the Oppenheimer-Snyder collapse [4]) was studied by Mann and Ross [5] who found that it – the three spacetime (3D) collapse – exhibits properties that entirely parallel those of the four spacetime (4D) Oppenheimer-Snyder collapse.

The quantum nature of the 4D Schwarzschild black hole was revealed in the paper by Ram, Ram and Ram [6] using a method in which one makes quantum mechanics from time-like geodesics, obtaining thereby a quantum equation which quantizes mass. The use of Bose's method [7] then reveals the quantum statistical nature of the 4D black hole, namely that it is a Bose-Einstein ensemble of quanta of mass equal to twice the Planck mass. In the present note we apply the method [6] to the 2+1 dimensional case.

The first integral of the time-like geodesics in three space-time dimensions with the BTZ line element (1) can be easily obtained, and is given by

$$\dot{r}^2 + L^2 \left(-\frac{8M}{r^2} + \frac{1}{\ell^2}\right) - 8M + \frac{r^2}{\ell^2} = E^2 \quad (2)$$

with E and L constants of integration. In Eq. (2) dot means differentiation with respect to the proper time. Following Ref. 6 we put $E = L = 0$ in (2), and obtain

$$\frac{1}{2}\dot{r}^2 + \frac{1}{2}\omega^2 r^2 = 4M \quad (3)$$

with $\omega = \frac{1}{\ell}$. Proper use of the Schrödinger prescription in (3) then leads to the quantum equation

$$\left(-\frac{1}{2r} \frac{\partial}{\partial r} \left(r \frac{\partial}{\partial r}\right) + \frac{1}{2}\omega^2 r^2\right) \psi = 4M\psi. \quad (4)$$

Thus the harmonic oscillator shows itself again, though in this case it is a two-dimensional one as opposed to the four-dimensional one in the 4D case.

With $U = r^{1/2}\psi$ and $M = \mu/4$, Eq. (4) takes the form

$$\left[-\frac{1}{2} \left(\frac{d^2}{dr^2} + \frac{1}{4r^2} \right) + \frac{1}{2} \omega^2 r^2 \right] U = \mu U \quad (5)$$

with ω equal to two [8,9]. The quantum equation (5) has the eigenvalues [10]

$$\mu_n = 2(n+1)\omega, \quad n = 0, 1, 2, \dots \quad (6)$$

Thus mass is quantized [11] as in (6). That is, the n th mass (μ) state is occupied by n pairs of mass quanta, each quantum being of $\omega = 2$ or of mass (energy) twice the Planck mass (energy); same as in the $4D$ case.

With the establishment of the connection of the energy element of the two-dimensional oscillator with the mass quantum of pure gravitation via Eqs. (3-6), one can again use Bose's method to calculate the entropy S of pure gravitation within a $3D$ black hole. Let the mass of the BTZ black hole as in the paper of Mann and Ross [5] be M_B and let it contain $2N$ quanta. Then $M_B = 2N\omega$. Let us rewrite Eq. (6) as

$$\epsilon_n = \frac{\mu_n}{2} = (n+1)\omega, \quad n = 0, 1, 2, \dots, \quad (7)$$

so that the n th ϵ -state contains n quanta. The thermodynamic quantity E in Bose's paper [7] is given by

$$E = \frac{M_B}{2} = N\omega, \quad (8)$$

same as in the $4D$ case. The entropy S is again calculable using the expression [6]

$$S = \frac{E}{T} - N \left(e^{\omega/T} - 1 \right) \ln \left(1 - e^{-\omega/T} \right), \quad (9)$$

or

$$S = \frac{M_B}{2T} - \frac{M_B}{2\omega} \left[\left(e^{\omega/T} - 1 \right) \ln \left(1 - e^{-\omega/T} \right) \right]. \quad (10)$$

For $\frac{\omega}{T} \ll 1$, i.e. for large M_B , Eq. (10) reduces to

$$S = \frac{M_B}{2T} \left[1 - \ln \left(\frac{\omega}{T} \right) \right]. \quad (11)$$

For the 2+1 dimensional black hole, if we take [2] $T = \frac{r_B^+}{2\pi\ell^2} = \frac{r_B^+\omega^2}{2\pi}$ with $r_B^{+2} = 8M_B\ell^2 = \frac{8M_B}{\omega^2}$, then Eq. (11) becomes

$$S = \frac{2\pi r_B^+}{4} \left[\frac{1 - \ell n\left(\frac{\omega}{T}\right)}{4} \right] = \frac{2\pi r_B^+}{4} \left[\frac{1 - \ell n\left(\frac{\pi}{\sqrt{2M_B}}\right)}{4} \right]. \quad (12)$$

Eq. (12) says that the entropy of a 3D black hole is given by not just $\frac{2\pi r_B^+}{4}$ but by $\frac{2\pi r_B^+}{4}$ times a monotonically increasing function of the black hole mass M_B [12,13]. This is the difference between the 2+1 dimensional black hole and the 3+1 dimensional black hole for which the entropy comes out to be $\frac{A}{4}$ (A being the area of the 4D black hole and $2\pi r_B^+$ being its 3D analog).

The above investigation demonstrates – plainly – that the method [6] originally given for the 4D case works as well for the 3-dimensional case. The investigation, moreover, brings out – unambiguously – the importance of quantum statistical effects.

Acknowledgements

The authors thank N. Ram for a critical reading of the manuscript. One of us (BR) thanks A. Ram, T. Halverson, R.S. Bhalerao, N. Banerjee and R. Tipton for useful conversations, and TIFR and IUCAA for pleasant stays.

References and Footnotes

- [1] M. Bañados, C. Teitelboim and J. Zanelli, Phys. Rev. Lett. 69, 1849 (1992).
- [2] S. Carlip, Class. Quant. Grav. 22, R85 (2005).
- [3] Whereas in Ref. 1 the units used are such that $8G = 1 = c$, we use units in which $G = 1 = c = \hbar = k$.
- [4] J.R. Oppenheimer and H. Snyder, Phys. Rev. 56, 455 (1939).
- [5] R.B. Mann and S.F. Ross, Phys. Rev. D47, 3319 (1993).
- [6] B. Ram, A. Ram, and N. Ram, arXiv: gr-qc/0504030; B. Ram, Phys. Lett. A265, 1 (2000).
- [7] S.N. Bose, Z. Phys. 26, 178 (1924); English translation in O. Theimer and B. Ram, Am. J. Phys. 44, 1056 (1976).
- [8] Since Eq. (5) with $\omega = 2$ is a map of [see Ref. 6]
$$\left[-\frac{1}{2} \left(\frac{d^2}{d\rho^2} + \frac{1}{4\rho^2} \right) - \frac{M}{\rho} \right] \phi(\rho) = -\frac{1}{2} \phi(\rho).$$
In this sense, M is analog of the Schwarzschild mass; see also Ref. 5.
- [9] V. Cardoso and J. Lemos take $\ell = 1$ in their paper [Phys. Rev. D63, 124015 (2001)] which corresponds to $\omega = 1$.
- [10] H.A. Mavromatis, Exercises in Quantum Mechanics (D. Reidel Publishing Co., Dordrecht, Holland, 1986), p. 95.
- [11] In this connection, see also M.R. Setare, Class. Quant. Grav. 21, 1453 (2004). We thank M. Setare for bringing this paper to our attention.
- [12] In this connection see p. 2869 of S. Carlip, Class. Quant. Grav. 12, 2853 (1995), and A. Ghosh and P. Mitra, Mod. Phys. Lett. A11, 1231 (1996).
- [13] For $\frac{\omega}{T} = 10^{-20}$ or $M_B = 5 \times 10^{40}$, the numerical value of $\left[\frac{1-\ell n(\omega/T)}{4} \right]$ is $\left[\frac{1+20\ell n 10}{4} \right]$, and the value increases monotonically by $\frac{\ell n 10}{4}$ as $\frac{\omega}{T}$ decreases by a factor of 10 or M_B increases by a factor of 100.